Patient Education

Clinical Center NATIONAL INSTITUTES OF HEALTE

Don't let food make you sick! Foods to Choose

Food safety tips for those at increased risk of food-borne illness

Food	Choose	Do not eat or drink
Fruits and vegetables	 Pasteurized juice Fresh fruits/vegetables that have no bruises, cuts or mold on them Frozen fruits/vegetables (except berries) Cooked, pasteurized, or canned berries Cooked vegetable sprouts Cooked or canned fruits or vegetables Well-scrubbed produce Use a vegetable brush for rough surfaces like melon skins. Produce washed right before consuming (to reduce risk of mold) Pre-washed, bagged greens that are well refrigerated, consumed before the expiration date and free of "off" odors and wilted leaves 	 Unpasteurized juices Pre-cut, damaged, or loose pieces of fruit (such as grapes off the stem) All raw and frozen berries (they are hard to clean and get moldy quickly) Raw bean sprouts Avoid when eating out: Buffet-style salad bars Deli salads such as fruit, potato, egg or pasta salad made with raw fruit or vegetables Salsas (unless pasteurized) Raw fruits or vegetables (since you can't be sure of safe food handling)

Food	Choose	Do not eat or drink
Nuts	 Commercially packaged nut butters (peanut, almond, cashew, soy) Commercially packaged, roasted nuts Almonds (all types are pasteurized now) 	 Freshly ground nut butters Nuts in the shell (because of mold risk) Unroasted nuts sold in bulk from bins
Meat, fish, poultry, soy	 Pasteurized or boiled tofu Meats cooked to these temperatures: (Insert thermometer into thickest part.) Poultry 165° F Ground meat (like hamburger) 160° F Steaks and roasts 145° F Egg Dishes 160° F Fish 145° F (firm flesh, flakes easily) Luncheon meats heated to 165° F Well-done meat cooked to order, served hot 	 Uncooked tofu (unless pasteurized) Raw or undercooked meat or fish, including: smoked salmon (lox), sushi, raw oysters, "Ahi" tuna, sashimi (raw fish) Avoid when eating out: Meat prepared: "Rare" "Seared" "Cured" Luncheon meats (unless heated to 165° F) Items kept warm

Food	Choose	Do not eat or drink
Milk, cheese, yogurt	 Pasteurized milk Cheese made from pasteurized milk Yogurt made from pasteurized milk Ask your doctor if you should avoid "live and active cultures" when your "counts" are low. 	 Raw milk or yogurt made with raw milk Cheese made from raw milk Cheese with herbs or other added items (such as dill, peppers) Cheese with mold (such as blue cheese) Unpasteurized cheese (Look for pasteurized versions of cheeses such as feta and brie.)
Eggs	 Well-cooked eggs (Hard-boiled, scrambled) Pasteurized shell eggs (Sold in cartons next to regular eggs; can be used to safely prepare softer-style eggs) Pasteurized liquid egg products (like low-cholesterol eggs, egg whites) 	 Eggs "over easy," soft poached, soft boiled, or "sunny side up" (unless pasteurized shell eggs are used) Raw eggs or products made with raw eggs such as salad dressings, protein shakes and raw dough. Use pasteurized egg products for these foods instead.

This information is prepared specifically for persons taking part in clinical research at the National Institutes of Health Clinical Center and may not apply to patients elsewhere. If you have questions about the information presented here, talk to a member of your health care team.

Products/resources named serve as examples and do not imply endorsement by NIH. The fact that a certain product/resource is not named does not imply that such product/resource is unsatisfactory.

National Institutes of Health Clinical Center Bethesda, MD 20892

Questions about the Clinical Center?

http://www.cc.nih.gov/comments.shtml